Milk Replacer currently stocked and available



PREMIUM
20/20 MOS
CALF MILK REPLACER
NON-MEDICATED

GUARANTEED ANALYSIS

Crude Protein, min.	20.00%
Crude Fat, min.	20.00%
Crude Fiber, max.	0.15%
Calcium, min.	0.75%
Calcium, max.	1.25%
Phosphorus, min.	0.60%
Vitamin A, min.	30,000 IU/LB
Vitamin D_3, min.	10,000 IU/LB
Vitamin E, min.	200 IU/LB

INGREDIENTS

Dried Whey, Dried Whey Protein Concentrate, Dried Whey Product, Animal and Vegetable Fat (Preserved with BHT), Dried Skimmed Milk, Coconut Oil, Lecithin, Dried Milk Protein, Polyoxyethylene Glycol (400) Mono and Dioleates, Primary Dried Yeast, Calcium Carbonate, Dicalcium Phosphate, Magnesium Sulfate, l-Lysine, Choline Chloride, Silicon Dioxide, dl-Methionine, Vitamin E Supplement, Iron Amino Acid Complex, Ascorbic Acid, Manganese Amino Acid Complex, Zinc Amino Acid Complex, Vitamin A Supplement, Niacinamide, Vitamin D Supplement, Copper Amino Acid Complex, Selenomethionine Hydroxy Analogue, Artificial Flavor, d-Calcium Pantothenate, Thiamine Mononitrate, Riboflavin, Vitamin B_{12} Supplement, Pyridoxine Hydrochloride, Biotin, Menadione Sodium Bisulfite Complex (Source of Vitamin K_3), Calcium Iodate, Folic Acid.

See Feeding Instructions on Reverse Side

MANUFACTURED BY
STRAUSS FEEDS, LLC
WATERTOWN, WI 53098

Net Weight 50 LB (22.68 KG)

2781101 1023



FEEDING INSTRUCTIONS

It is recommended that calves be given 2 to 3 quarts of good quality colostrum as soon as possible after birth, and another 2 to 3 quarts within the first 24 hours of life. Second milking colostrum may be fed afterwards for up to 3 days. Starting on the 4th day, feed a high quality replacer according to the following directions.

MIXING

Into 2 quarts of warm (110°F) clean water, add 10 ounces (1 cup)* of milk replacer powder and mix thoroughly. For larger batches, mix 1 1/4 pounds of milk replacer powder per gallon of warm water.

FEEDING

Feed the reconstituted milk replacer at a rate of 2 quarts per 100 pounds of body weight twice daily (4 quarts per 100 pounds of body weight per day). Large breed calves generally require 2 to 2-1/2 quarts per feeding twice daily, while smaller breed calves may need 1-1/2 to 2 quarts twice daily.

MANAGEMENT SUGGESTIONS

- For best mixing results, sprinkle milk replacer powder onto water.
- Use of a wire whisk or mechanical agitation provides good mixing.
- Feed reconstituted milk replacer shortly after mixing.
- Feeding and mixing equipment should be cleaned and sanitized after each use.
- Clean, fresh water should be available at all times.
- Calves should have access to a quality texturized feed after 4 days of age.
- Hay should not be offered until calves are weaned.
- Wean calves when they are consuming 1-1/2 pounds of starter feed per day for 3 consecutive days.

*The measuring cup provided in the bag holds approximately 10 ounces of milk replacer powder. For greater accuracy, a suitable scale is recommended.

**** Please call the office for any questions****

****Additional Milk replacers can be specially ordered upon request***

BROWN'S®

FEEDING GENERATIONS OF FARMS SINCE 1871

127 South Furnace Street Birdsboro PA 19508



610-582-2741

BrownsFeeds.com